

Closure, Properties and Closure Properties of Multirelations

Rudolf Berghammer

Christian-Albrechts-Universität zu Kiel

Walter Guttmann

University of Canterbury

1. Multirelations
2. Reflexive-Transitive Closure
3. Properties and their Closure
4. Topological Contact

Context and Method

- multirelations in program semantics, games, topological contact
- systematically investigate their properties

- express multirelational operations using relations
- study properties of operations
- abstract properties to weak algebras
- derive theory in these algebras

Relations and Multirelations

- state space $A = \{1, 2, 3\}$
- relation $\subseteq A \times A$

	1	2	3
1	■	■	□
2	□	□	□
3	□	■	□

multirelation $\subseteq A \times 2^A$

	\emptyset	3	2	23	1	13	12	123
1	□	□	□	■	■	□	■	□
2	□	□	□	□	□	□	□	□
3	■	□	□	□	□	□	□	□

- Boolean algebra with \cup , \cap , $^-$
- composition
- converse \cdot^c , dual \cdot^d

Multirelational Constants

O \equiv

	\emptyset	3	2	23	1	13	12	123
1								
2								
3								

E \equiv

	\emptyset	3	2	23	1	13	12	123
1					■	■	■	■
2			■	■			■	■
3		■		■		■		■

T \equiv

	\emptyset	3	2	23	1	13	12	123
1	■	■	■	■	■	■	■	■
2	■	■	■	■	■	■	■	■
3	■	■	■	■	■	■	■	■

U \equiv

	\emptyset	3	2	23	1	13	12	123
1					■			
2			■					
3		■						

Relational Composition

	1	2	3
1	■	■	□
2	□	□	□
3	□	■	□

	1	2	3
1	□	□	■
2	□	■	□
3	■	■	■

	1	2	3
1	□	■	■
2	□	□	□
3	□	■	□

$$(QR)_{x,z} \Leftrightarrow \exists y \in A : Q_{x,y} \wedge R_{y,z}$$

Multirelational Composition

	\emptyset	3	2	23	1	13	12	123
1								
2								
3								

	\emptyset	3	2	23	1	13	12	123
1								
2								
3								

	\emptyset	3	2	23	1	13	12	123
1								
2								
3								

$$(Q; R)_{x,z} \Leftrightarrow \exists Y \in 2^A : Q_{x,Y} \wedge \forall y \in Y : R_{y,z}$$

Up-closed Multirelations

not up-closed

	\emptyset	3	2	23	1	13	12	123
1				■	■		■	
2								
3	■							

up-closed

	\emptyset	3	2	23	1	13	12	123
1				■	■	■	■	■
2								
3	■	■	■	■	■	■	■	■

$$\forall x \in A : \forall Y, Z \in 2^A : R_{x,Y} \wedge Y \subseteq Z \Rightarrow R_{x,Z}$$

Relational Operations for Multirelations

right residual $Q \setminus R = \overline{Q^c R}$

symmetric quotient $Q \div R = (Q \setminus R) \cap (R \setminus Q)^c$

subset relation : $2^A \leftrightarrow 2^A$ $S = E \setminus E$

multirelational composition $Q ; R = Q(E \setminus R)$

R up-closed if $R = RS$

Unit and Zero of Multirelations

left unit $E;R = E(E \setminus R) = R$

right unit $R;E = R(E \setminus E) = RS = R$ if R up-closed

left zero $O;R = O$

$T;R = T$

Laws of Multirelations

all multirelations

$$O;R = O$$

$$E;R = R$$

$$T;R = T$$

$$R;E \supseteq R$$

$$Q \subseteq R \Rightarrow P;Q \subseteq P;R$$

$$(P \cup Q);R = P;R \cup Q;R$$

$$(P \cap Q);R \subseteq P;R \cap Q;R$$

$$(P;Q);R \subseteq P;(Q;R)$$

up-closed multirelations

$$R;E = R$$

$$(P \cap Q);R = P;R \cap Q;R$$

$$(P;Q);R = P;(Q;R)$$

Algebraic Structures

bounded join-semilattice

$$\begin{aligned}x + (y + z) &= (x + y) + z \\x + y &= y + x\end{aligned}$$

$$\begin{aligned}x + x &= x \\0 + x &= x\end{aligned}$$

pre-left semiring

$$\begin{aligned}(x \cdot y) + (x \cdot z) &\leq x \cdot (y + z) \\(x \cdot z) + (y \cdot z) &= (x + y) \cdot z \\0 &= 0 \cdot x\end{aligned}$$

$$\begin{aligned}(x \cdot y) \cdot z &\leq x \cdot (y \cdot z) \\x &\leq x \cdot 1 \\x &= 1 \cdot x\end{aligned}$$

left residual

$$x \cdot y \leq z \Leftrightarrow x \leq z/y$$

Reflexive-Transitive Closure

recursion modelled by

$$f(x) = 1 + x \cdot y$$

$$g(x) = 1 + y \cdot x$$

$$h(x) = 1 + y + x \cdot x$$

least prefixpoint

$$f(\mu f) \leq \mu f \quad f(x) \leq x \Rightarrow \mu f \leq x$$

if μf , μg , μh exist then

$$\mu f \leq \mu g = \mu h$$

Properties of Multirelations

up-closed $R;E = R$

total $R;T = T$

co-total $R;O = O$

\cup -distributive $R;(P \cup Q) = R;P \cup R;Q$

\cap -distributive $R;(P \cap Q) = R;P \cap R;Q$

reflexive $E \subseteq R$

transitive $R;R \subseteq R$

idempotent $R;R = R$

co-reflexive $R \subseteq E$

dense $R \subseteq R;R$

contact $R;R \cup E = R$

test $R;T \cap E = R$

vector $R;T = R$

kernel $R;R \cap E = R;E$

co-test $R;O \cup E = R$

Algebraic Structures

$(S, +, \wedge, 0, \top)$ bounded distributive lattice,


$(S, +, \cdot, 0, 1)$ pre-left semiring and

$$\begin{aligned}\top &= \top \cdot x \\ x \cdot (y \cdot z) &= (x \cdot (y \cdot 1)) \cdot z \\ (x \cdot z) \wedge (y \cdot z) &= ((x \cdot 1) \wedge (y \cdot 1)) \cdot z\end{aligned}$$

dual

$$\begin{aligned}(x \cdot y)^d &= (x \cdot 1)^d \cdot y^d \\ (x + y)^d &= x^d \wedge y^d \\ x^{dd} &= x \\ 1^d &= 1\end{aligned}$$

Relationships between Properties



Closure Properties

	O	E	T	U	\cap	;	^d
total	—	■	■	■	□	□	▽
co-total	■	■	—	■	■	■	▲
transitive	■	■	■	—	■	—	▼
dense	■	■	■	■	—	—	△
reflexive	—	■	■	■	■	■	▼
co-reflexive	■	■	—	■	■	■	▲
idempotent	■	■	■	—	—	—	□
up-closed	■	■	■	■	■	■	■
\cup -distributive	■	■	■	■	—	□	▽
\cap -distributive	■	■	■	—	□	□	△
a contact	—	■	■	—	■	—	▼
a kernel	■	■	—	■	—	—	▲
a \cup -distributive contact	—	■	■	—	—	—	▼
a \cap -distributive kernel	■	■	—	—	—	—	▲
a test	■	■	—	■	■	■	▼
a co-test	—	■	■	■	■	■	▲
a vector	■	—	■	■	■	■	■

Topological Contact

- according to G. Aumann (1970)
- set of persons A
- set of topics T
- $t(x)$ = topics person x is interested in

$$t : A \rightarrow 2^T$$

- contact multirelation $R : A \leftrightarrow 2^A$

$$R_{x,Y} \Leftrightarrow t(x) \subseteq \bigcup_{y \in Y} t(y)$$

Axioms of Contact Relations

$$(K_0) \quad \neg \exists x \in A : R_{x, \emptyset}$$

$$(K_1) \quad \forall x \in A : R_{x, \{x\}}$$

$$(K_2) \quad \forall x \in A : \forall Y, Z \in 2^A : R_{x, Y} \wedge Y \subseteq Z \Rightarrow R_{x, Z}$$

$$(K_3) \quad \forall x \in A : \forall Y, Z \in 2^A : R_{x, Y} \wedge (\forall y \in Y : R_{y, Z}) \Rightarrow R_{x, Z}$$

$$(K_4) \quad \forall x \in A : \forall Y, Z \in 2^A : R_{x, Y \cup Z} \Leftrightarrow R_{x, Y} \vee R_{x, Z}$$

(K_1) – (K_3) contact relation

(K_0) – (K_4) topological contact relation

Examples of Topological Contact

- \in $A \leftrightarrow 2^A$
- $R_{x,Y} \Leftrightarrow \exists y \in Y : f(x) = f(y)$ where $f : A \rightarrow B$
- $R_{x,Y} \Leftrightarrow \exists y \in Y : x \leq y$ $\mathbb{N} \leftrightarrow 2^{\mathbb{N}}$
- $R_{x,Y} \Leftrightarrow \exists y_1, y_2 \in Y : y_1 \leq x \leq y_2$
- $R_{x,Y} \Leftrightarrow \exists y_i \in Y : \exists r_i \in \mathbb{Q} : x = \sum r_i y_i$ $\mathbb{R}^n \leftrightarrow 2^{\mathbb{R}^n}$
- $R_{x,Y} \Leftrightarrow \exists y_i \in Y : \exists r_i \in \mathbb{Q}_0^+ : \sum r_i = 1 \wedge x = \sum r_i y_i$
- $R_{x,Y} \Leftrightarrow \forall \varepsilon > 0 : \exists y \in Y : d(x, y) < \varepsilon$

satisfy (K_0) – (K_3) , some also (K_4)

Axioms using Multirelational Operations

(K_0)	$R;O = O$	co-total
(K_1)	$E \subseteq R$ (if R up-closed)	reflexive
(K_2)	$R;E = R$	up-closed
(K_3)	$R;R \subseteq R$	transitive
(K_4)	$R;(P \cup Q) = R;P \cup R;Q$ (if R up-closed)	\cup -distributive

Conclusion

- multirelations describe topological contact
- also consider not up-closed multirelations
- many results hold in weak algebras

- study connections to topology and closure systems
- generate further counterexamples
- give complete axioms